Calculus refresher Statistical Natural Language Processing 1

Çağrı Çöltekin

University of Tübingen Seminar für Sprachwissenschaft

Winter Semester 2024/2025

version: 4a58e24 @2024-12-03

Calculus and NLP

- *•* Our main use case is finding minima:
	- **–** Supervised ML models are (typically) trained by minimizing the error, or a *loss function*
	- **–** Differential calculus allows efficiently searching minima, and determining where minima are
- *•* Integrals will also be handy for calculating probabilities

Today's plan

Very brief introductions to

- *•* Limits
	- **–** We are mainly interested in for defining derivatives
- *•* Derivatives
	- **–** Are the central topic for us: training an ML system relies on derivation (but there are also other interesting uses)
- *•* Integrals
	- **–** Mainly for probability theory, but without integrals, the derivatives would also be incomplete

Limits

- *•* Limit is the value that a function approaches as its argument approaches (arbitrarily close, but not equal) to some value.
- *•* We write

$$
\lim_{x\to c}f(x)
$$

for the value of function f as x approaches to c.

• If the value of the function at $x = c$ is a number, the limit is $f(c)$

$$
\lim_{x\to 2} x^2 = 2^2 = 4
$$

- *•* If the result is [∞], the limit does not exist
- *•* Interesting cases are when the function is discontinuous, or undefined, e.g., f(x) is $0/0$, ∞/∞ , $\infty - \infty$

Example (1)

$$
\lim_{x\to 1}\frac{1}{x}=
$$

Example (1)

$$
\lim_{x \to 1} \frac{1}{x} = 1
$$

$$
\lim_{x\to 0}\frac{1}{x}=
$$

Example (1)

$$
\lim_{x \to 1} \frac{1}{x} = 1
$$

lim $x \rightarrow 0$ 1 $\frac{1}{x} = \pm \infty$

Example (1)

$$
\lim_{x\to 1}\frac{1}{x}=1
$$

$$
\lim_{x\to 0}\frac{1}{x}=\,\pm\,\infty
$$

More precisely,

$$
\lim_{x \to 0^{-}} \frac{1}{x} = -\infty
$$

$$
\lim_{x \to 0^{+}} \frac{1}{x} = \infty
$$

Example (2)

$$
f(x) = \begin{cases} -1 & x < 0 \\ 1 & x > 0 \end{cases}
$$

$$
\lim_{x\to 0} f(x) = ?
$$

Example (2)

$$
f(x) = \begin{cases} -1 & x < 0\\ 1 & x > 0 \end{cases}
$$

$$
\lim_{x \to 0} f(x) = ?
$$

$$
\lim_{x \to 0^{-}} f(x) = -1
$$

$$
\lim_{x \to 0^{+}} f(x) = 1
$$
limit does not exist

Example (3)

$$
\lim_{x \to 4} \frac{x - 4}{\sqrt{x} - 2}
$$

Example (3)

$$
\lim_{x \to 4} \frac{x-4}{\sqrt{x}-2}
$$

$$
\lim_{x \to 4^{-}} \frac{x-4}{\sqrt{x}-2} = 4
$$

$$
\lim_{x \to 4^{+}} \frac{x-4}{\sqrt{x}-2} = 4
$$

 $x \rightarrow 4^+$

Rules for limits

- $\lim_{x \to c} (f(x) + g(x)) = \lim_{x \to c} f(x) + \lim_{x \to c} g(x)$
- $\lim_{x \to c} af(x) = a \lim_{x \to c} f(x)$
- $\lim_{x \to c} (f(x)g(x)) = \lim_{x \to c} f(x) \lim_{x \to c} g(x)$

Derivatives

- *•* Derivative of a function f(x) is another function f *′* (x) indicating the rate of change in $f(x)$
- Alternatively: $f'(x) = \frac{df}{dx}(x)$
- *•* When derivative exists, it determines the tangent line to the function at a given point
- *•* Example from physics: velocity is the derivative of the position
- *•* Our main interest:
	- **–** the points where the derivative is 0 are the stationary points (maxima, minima, inflection points)
	- **–** the derivative evaluated at other points indicate the direction and steepness of the curve defined by the function

Example: derivatives

- $f'(x)$ is negative when $f(x)$ is decreasing, positive when it is increasing
- *•* The absolute value of f *′* (x) indicates how fast $f(x)$ changes when x changes
- *•* f *′* (x) = 0 when at a *stationary point*
- *•* f *′* (a) is a (good) approximation to the $f(x)$ near the α

Derivative of a function

definition

$$
f'(x) = \lim_{\Delta \to 0} \frac{f(x + \Delta) - f(x)}{\Delta}
$$

Example: calculating derivatives using the definition

$$
\frac{d}{dx}x^2 = \lim_{\Delta \to 0} \frac{(x + \Delta)^2 - x^2}{\Delta}
$$

$$
= \lim_{\Delta \to 0} \frac{x^2 + 2\Delta x + \Delta^2 - x^2}{\Delta}
$$

$$
= 2x
$$

Some derivatives to know

- Powers: $\frac{d}{dx}x^n = nx^{n-1}$
- Trigonometric functions:
 $\frac{d}{dx} \sin(x) = \cos(x)$
 $\frac{d}{dx} \cos(x) = -\sin(x)$

$$
\frac{\frac{d}{dx}\cos(x)}{\frac{d}{dx}\tan(x)} = 1 + \tan^2(x)
$$

- Powers of $e: \frac{d}{dx}e^x = e^x$
- Natural logarithm: $\frac{d}{dx} \ln(x) = \frac{1}{x}$

General rules for derivatives

• Sum rule:

$$
\frac{d}{dx}f(x) + g(x) = \frac{d}{dx}f(x) + \frac{d}{dx}g(x)
$$

• Product rule:

$$
\frac{d}{dx}f(x)g(x) = g(x)\frac{d}{dx}f(x) + f(x)\frac{d}{dx}g(x)
$$

• Quotient rule:

$$
\frac{d}{dx}\frac{f(x)}{g(x)} = \frac{g(x)\frac{d}{dx}f(x) - f(x)\frac{d}{dx}g(x)}{g^2(x)}
$$

• Chain rule: if $f(x) = h(g(x))$

$$
\frac{df}{dx} = \frac{dh}{dg} \frac{dg}{dx}
$$
, or $f'(x) = h'(g(x))g'(x)$

Cain rule: examples

d $\frac{u}{dx}e^x$

 $\mathbf d$ $rac{a}{dx} 2^{\sin x}$

Derivatives and extrema

- *•* Derivative of a function is 0 at minimum, maximum and inflection points
- *•* Derivative is useful for optimization (minimization of maximization) problems
- *•* We need additional tests to determine the type of critical points

Derivatives and extrema

- *•* Derivative of a function is 0 at minimum, maximum and inflection points
- *•* Derivative is useful for optimization (minimization of maximization) problems
- *•* We need additional tests to determine the type of critical points

Higher order derivatives

- *•* Higher order derivatives, particularly second derivative, are useful in many applications
	- **–** Determining the type of critical points
	- **–** Polynomial approximations to functions
- *•* Notation:
	- Second derivative: $f''(x) = \frac{d^2 f}{dx^2}$
	- **–** nth derivative: $f^{(n)}(x) = \frac{d^n \vec{f}}{dx^n}$

Second derivatives and extrema

- *•* Second derivatives are useful for determining the type of critical points
	- **–** f *′′*(x) < 0 if f(x) is concave down (*∩*)
	- **–** f *′′*(x) > 0 if f(x) is concave up (*∪*) **–** f *′′*(x) = 0 if f(x) is flat

Differentiable functions and continuity

- *•* A function is said to be *differentiable* if its derivative exists in every point in its domain
- *•* This concept is important when we wan to use optimization techniques based on derivatives
- *•* A differentiable function is also continuous, but a continuous function is not necessarily differentiable

Differentiability

Are these functions differentiable?

Partial derivatives and gradient

- *•* In ML, we are often interested in (error) functions of many variables
- *•* A partial derivative is derivative of a multivariate function with respect to a single variable, noted $\frac{\partial f}{\partial x}$
- *•* A very useful quantity, called *gradient*, is the vector of partial derivatives with respect to each variable

$$
\nabla f(x_1,\ldots,x_n) = \left(\frac{\partial f}{\partial x_1},\ldots,\frac{\partial f}{\partial x_n}\right)
$$

- *•* Gradient points to the direction of the steepest change
- Example: if $f(x, y) = x^3 + yx$

$$
\nabla f(x,y) = \left(3x^2 + y, x\right)
$$

Gradient visualization

Integrals

- *•* Integral is the reverse of the derivative (anti-derivative)
- The indefinite integral of $f(x)$ is noted $F(x) = \int f(x) dx$
- *•* We are often interested in definite integrals

$$
\int_{\alpha}^{b}f(x)dx=F(b)-F(\alpha).
$$

• Integral gives the area under the curve

Numeric integrals & infinite sums

- *•* As the width of the rectangles converges to 0 (or number of rectangles becomes ∞), the sum converges to the area under the curve
- *•* Integration is 'infinite summation'
- *•* When integration is not possible with analytic methods, we resort to numeric integration

Summary / next

We reviewed three main concepts from calculus

- *•* Limits
- *•* Derivatives
- *•* Integrals

Summary / next

We reviewed three main concepts from calculus

- *•* Limits
- *•* Derivatives
- *•* Integrals

Next:

- *•* Regression again: through gradient optimization
- *•* Introduction to probability theory

Further reading

- *•* A nice video series: https://www.youtube.com/playlist?list= PLZHQObOWTQDMsr9K-rj53DwVRMYO3t5Yr
- *•* No concrete reading suggestions, but check https://www.openculture.com/free-math-textbooks